

Culture media on multiplates

| Ready multisection plates | Ref.No. | Packaging | Type |
|---|---------|-----------|---------|
| Bi-plates | | | |
| Brain Heart Infusion Streptomycin Agar / Brain Heart Infusion Gentamycin Agar For the selective isolation of <i>Enterococcus spp.</i> with high-level aminoglycoside resistance. | 2046 | plate | 90 mm/2 |
| Brain Heart Infusion Vancomycin Agar / Brain Heart Infusion Agar For the selective isolation of <i>Enterococcus spp.</i> with high-level vancomycin resistance. | 2045 | plate | 90 mm/2 |
| Buffered Charcoal Yeast Extract Agar (BCYE) / Buffered Charcoal Yeast Extract Agar (BCYE) with cysteine For the isolation of <i>Legionella spp.</i> | 2067 | plate | 90 mm/2 |
| Buffered Charcoal Yeast Extract Agar (BCYE) with vancomycin and colistin / Buffered Charcoal Yeast Extract (BCYE) Agar without cysteine with pimaricin For the isolation of <i>Legionella spp.</i> | 2071 | plate | 90 mm/2 |
| Bordetella Agar with 15% sheep blood / Bordetella Agar with charcoal and 7% horse blood For the cultivation of <i>Bordetella pertussis</i> and <i>Bordetella parapertussis</i> from clinical samples. | 2008 | plate | 90 mm/2 |
| Brucella Agar with 5% sheep blood, hemine and vit. K / CDC Kanamycin Vancomycin Agar with lysed sheep blood For the isolation of a wide variety of anaerobic microorganisms. | 2006 | plate | 90 mm/2 |
| CHROMagar VRE / CHROMagar KPC For the detection of antibiotic resistance mechanisms in bacteria. | 2019 | plate | 90 mm/2 |
| Campylobacter Agar (acc. Butzler) with 7% horse blood / Clostridium Difficile Agar with 7% sheep blood For the isolation of anaerobic bacteria from clinical samples. | 2007 | plate | 90 mm/2 |
| CHROMagar Salmonella / CHROMagar O157 For the isolation of <i>Salmonella spp.</i> and <i>Escherichia coli O157</i> from faecal specimens. | 2056 | plate | 90 mm/2 |
| CHROMagar ESBL / CHROMagar VRE For the detection of antibiotic resistance mechanisms in bacteria. | 2057 | plate | 90 mm/2 |
| CHROMagar Listeria / CHROMagar StrepB For isolation of <i>Streptococcus agalactiae</i> (GBS) and <i>Listeria spp.</i> from vaginal swabs. | 2062 | plate | 90 mm/2 |

| | | | |
|---|-------------|-------|---------|
| CHROMagar MRSA / Mannitol Salt Agar For the selective isolation of methicillin-resistant <i>Staphylococcus aureus</i> (MRSA). | 2026 | plate | 90 mm/2 |
| Columbia Agar with 5% sheep blood / Columbia Agar with 5% sheep blood For the cultivation of a wide variety of fastidious microorganisms. | 2001 | plate | 90 mm/2 |
| Columbia Agar with 5% sheep blood / Chocolate Agar For the cultivation of a wide variety of fastidious microorganisms. | 2061 | plate | 90 mm/2 |
| Columbia Agar with 5% sheep blood / Sabouraud Dextrose Chloramphenicol Agar For the isolation of fungi from faecal specimens. | 2065 | plate | 90 mm/2 |
| Columbia CNA (Colistin, Nalidixic Acid) Agar with 5% sheep blood / Chocolate Agar For the cultivation of a wide variety of fastidious microorganisms. | 2052 | plate | 90 mm/2 |
| Columbia CNA (Colistin, Nalidixic Acid) Agar with 5% sheep blood / Enterococcosel Agar For the isolation of gram positive microorganisms. | 2022 | plate | 90 mm/2 |
| Columbia CNA (Colistin, Nalidixic Acid) Agar with 5% sheep blood / Mannitol Salt Agar (MSA) For the isolation of gram positive microorganisms, especially <i>Staphylococcus aureus</i> . | 2032 | plate | 90 mm/2 |
| Columbia CNA (Colistin, Nalidixic Acid) Agar with 5% sheep blood / CHROMagar Orientation Agar For the isolation and enumeration of microorganisms in urine. | 2044 | plate | 90 mm/2 |
| Columbia CNA (Colistin, Nalidixic Acid) Agar with 5% sheep blood / Sabouraud Dextrose Chloramphenicol Gentamycin Agar For the isolation of gram positive bacteria and yeasts from clinical samples. | 2015 | plate | 90 mm/2 |
| Enterococcosel Agar / Arabinose Agar For the isolation and differentiation of <i>Enterococcus spp.</i> | 2059 | plate | 90 mm/2 |
| Enterococcosel Agar / Enterococcosel Vancomycin Agar For the selective isolation of <i>Enterococcus spp.</i> with high-level vancomycin resistance. | 2004 | plate | 90 mm/2 |
| Levine Eosin Methylene Blue Agar (LEMB) / Blood Agar with 5% sheep blood For the isolation and differentiation of gram negative enteric bacilli from faecal specimens. | 2043 | plate | 90 mm/2 |
| Levine Eosin Methylene Blue Agar (LEMB) / Mannitol Salt Agar (MSA) For the isolation of microorganisms from faecal specimens. | 2023 | plate | 90 mm/2 |
| MacConkey Agar with crystal violet / Blood Agar with 5% sheep blood For the isolation of microorganisms in urine. | 2100 | plate | 90 mm/2 |
| MacConkey Agar with crystal violet / CLED Agar (Cystine Lactose Electrolyte Deficient Agar) For the isolation of microorganisms in urine. | 2070 | plate | 90 mm/2 |

| | | | |
|--|-------------|-------|---------|
| MacConkey Agar with crystal violet / Columbia Agar with 5% sheep blood For the isolation of microorganisms in urine. | 2040 | plate | 90 mm/2 |
| MacConkey Agar with crystal violet / Columbia CNA (Colistin, Nalidixic Acid) Agar with 5% sheep blood For the isolation of microorganisms in urine. | 2050 | plate | 90 mm/2 |
| MacConkey Agar with crystal violet / Levine Eosin Methylene Blue Agar For the isolation of microorganisms from faecal specimens. | 2060 | plate | 90 mm/2 |
| MacConkey Agar with crystal violet / Sabouraud Dextrose Chloramphenicol Agar For the isolation of gram negative bacteria and yeasts. | 2010 | plate | 90 mm/2 |
| MacConkey Agar with crystal violet / Sabouraud Dextrose Chloramphenicol Gentamycin Agar For the isolation of gram negative bacteria and yeasts. | 2020 | plate | 90 mm/2 |
| MacConkey Sorbitol Agar / Yersinia CIN (Cefsulodin, Irgasan, Novobiocin) For the isolation and differentiation of enteric pathogens. | 2003 | plate | 90 mm/2 |
| Mannitol Salt Agar (MSA) / Edwards Agar with 5% sheep blood For the isolation of gram positive microorganisms. | 2049 | plate | 90 mm/2 |
| Oxacillin Resistance Staphylococci Isolation Medium (O.R.S.I.M) / Mannitol Salt Agar (MSA) For the selective isolation of methicillin-resistant <i>Staphylococcus aureus</i> . | 2069 | plate | 90 mm/2 |
| Oxacillin Resistance Staphylococci Isolation Medium (O.R.S.I.M) / Oxacillin Resistance Staphylococci Isolation Medium (O.R.S.I.M) with supplements For the selective isolation of methicillin-resistant <i>Staphylococcus aureus</i> . | 2095 | plate | 90 mm/2 |
| Phenylethyl Alcohol Blood Agar (PEA) / Schaedler KanaVanco (Kanamycin, Vancomycin) Agar with 5% sheep blood For the isolation of gram positive cocci and gram negative anaerobic bacteria. | 2039 | plate | 90 mm/2 |
| Sabouraud Dextrose Agar / CHROMagar Candida For the isolation of yeasts from vaginal swabs. | 2030 | plate | 90 mm/2 |
| Sabouraud Dextrose Chloramphenicol Agar / Fungisel Agar with phenol red For the isolation of dermatophytic fungi. | 2017 | plate | 90 mm/2 |
| Sabouraud Dextrose Agar / Sabouraud Dextrose Actidione Agar For the isolation of dermatophytic fungi and yeasts. | 2051 | plate | 90 mm/2 |
| Salmonella Shigella Agar / Bismuth Sulphite Agar (Wilson Blair) For the isolation of <i>Salmonella</i> spp. from faecal specimens. | 2037 | plate | 90 mm/2 |
| Salmonella Shigella Agar / Hektoen Enteric Agar For the isolation and differentiation of enteric pathogens. | 2014 | plate | 90 mm/2 |
| Schaedler Agar with 5% sheep blood / Schaedler KanaVanco (Kanamycin, Vancomycin) Agar with 5% sheep blood | 2021 | plate | 90 mm/2 |

For the isolation of gram negative anaerobic microorganisms.

Scheadler Agar with 5% sheep blood, hemine and vit. K / Schaedler KanaVanco (Kanamycin, Vancomycin) Agar with 5% horse blood 2072 plate 90 mm/2
For the isolation of gram negative anaerobic microorganisms.

Salmonella Shigella Agar / Xylose Lysine Desoxycholate Agar (XLD Agar) 2025 plate 90 mm/2
For the isolation and differentiation of enteric pathogens from faecal specimens.

Scheadler Agar with 5% sheep blood / Scheadler NeoVanco (Neomycin, Vancomycin) Agar with 5% sheep blood 2038 plate 90 mm/2
For the isolation of gram positive anaerobic cocci.

Tryptic Soy Agar with NAD (factor V) / Tryptic Soy Agar with NAD and hemine 2005 plate 90 mm/2
For the isolation and differentiation of *Haemophilus* spp.

Tri-plates*

Brain Heart Infusion Agar (BHI) with vancomycin (6 mg/l) / Brain Heart Infusion (BHI) with streptomycin (2000 mg/l) / Brain Heart Infusion (BHI) with gentamycin (500 mg/l) 2058 plate 90 mm/3
For the selective isolation of *Enterococcus* spp. with high-level aminoglycoside resistance.

MacConkey Agar with crystal violet / Mannitol Salt Agar (MSA) / Azide Blood with 5% sheep blood 2013 plate 90 mm/3
For the isolation and differentiation of gram positive cocci and gram negative microorganisms from throat swabs.

MacConkey Agar / Mannitol Salt Agar / Columbia Agar with 5% sheep 2054 plate 90 mm/3
For the isolation and differentiation of gram positive cocci and gram negative microorganisms from clinical samples.

MacConkey Agar / Mannitol Salt Agar / Edwards Agar with 5% sheep blood 2053 plate 90 mm/3
For the isolation of microorganisms including *Streptococcus agalactiae*.

Tetra-plates*

Brain Heart Infusion Agar (BHI) with vancomycin (6 mg/l) / Brain Heart Infusion Agar (BHI) with streptomycin (2000 mg/l) / Brain Heart Infusion Agar (BHI) with gentamycin (500 mg/l) / Brain Heart Infusion (BHI) Agar 2064 plate 90 mm/4
For the selective isolation of *Enterococcus* spp. with high-level aminoglycoside and glycopeptide resistance.

MacConkey Agar / Mannitol Salt Agar / Columbia Agar with 5% sheep blood / Chocolate Agar 2055 plate 90 mm/4
For the isolation and differentiation of gram positive and Gram negative microorganisms from clinical samples.

- **available on request**